

# We ask

## Gombe State Ministry of Health should ensure:

- ✓ Regular and adequate provision of Intermittent Preventive Treatment for pregnant women
- ✓ Timely distribution of data tools to all PHC facilities to address under reporting of data especially ARV prophylaxis
- ✓ Timely release of budgeted funds for HMIS to facilitate PHC data quality and entry into the DHIS2
- ✓ Production and timely distribution of IEC materials that give information to health workers to support and encourage early ANC visits to the health facilities by pregnant women
- ✓ Airing of radio, Jingles and TV programmes to raise awareness on acceptance of early ANC for pregnant women



[mamaye.org.ng](http://mamaye.org.ng)

A detailed reference sheet that explains how these figures are calculated, national and international standards and any limitations in the data is available at:  
[http://mamaye.org/sites/default/files/docs/gombe\\_MNCH\\_reference\\_sheet2018.pdf](http://mamaye.org/sites/default/files/docs/gombe_MNCH_reference_sheet2018.pdf)

Produced by the Gombe State Accountability Mechanism for Maternal and Newborn Health (GoSAM) in September 2018, with support from MamaYe-Evidence for Action



# MATERNAL, NEWBORN AND CHILD HEALTH (MNCH) SCORECARD

## GOMBE STATE, NIGERIA

January – June 2018



LOCAL GOVERNMENT AREAS

INDICATORS	STATE SUMMARY	LOCAL GOVERNMENT AREAS										
		AKKO	BALANGA	BILIRI	DUKKU	FUNAKAYE	GOMBE	KATUNGO	KWAMI	NAFADA	SHONGOM	YAMALTU
% of pregnant women reporting for antenatal care before 20 weeks	30%	27%	43%	32%	20%	24%	30%	34%	29%	30%	42%	27%
% of pregnant women that attended at least 4th ANC1	34%	22%	38%	42%	28%	33%	34%	40%	24%	26%	40%	51%
% of pregnant women who received two doses of Intermittent Preventive Treatment for Malaria	30%	27%	17%	34%	38%	34%	42%	26%	21%	27%	37%	15%
% of pregnant women who attended post natal clinic visit within 3 days delivery	14%	12%	6%	19%	6%	19%	15%	18%	20%	17%	25%	20%
% of HIV positive pregnant women on antiretroviral therapy (ARV) prophylaxis <sup>1</sup>	6%	1%	7%	6%	450%	1%	11%	17%	0%	0%	3%	7%
% of Deliveries taken by a skilled birth attendant	79%	89%	44%	85%	87%	61%	95%	93%	97%	89%	83%	77%
% of fully immunized children <1 year	20%	18%	25%	9%	15%	27%	34%	17%	4%	7%	34%	21%
% of children 0-6 months exclusively breastfed	21%	11%	34%	34%	4%	29%	52%	17%	5%	9%	21%	11%
% of new diarrhoea cases in under five year olds who are given ORS&zinc supplementation	56%	38%	24%	76%	50%	66%	82%	58%	89%	52%	50%	46%
% of new family planning acceptors (among all females using any method of contraception)	53%	51%	66%	60%	62%	37%	45%	63%	47%	71%	68%	52%
% of females aged 15 - 49 years using any method of modern contraception	6%	5%	4%	5%	4%	4%	16%	8%	3%	2%	7%	7%
% of births which are stillborn <sup>2</sup>	3%	2%	2%	1%	5%	2%	4%	1%	3%	3%	1%	3%
% of caesarian deliveries <sup>3</sup>	3%	1%	1%	3%	2%	2%	8%	3%	0%	0%	0%	1%
% of assisted vaginal delivery <sup>4</sup>	5%	5%	3%	9%	10%	10%	1%	2%	1%	7%	17%	7%
% of deliveries monitored by Partograph	45%	33%	13%	53%	52%	32%	82%	18%	56%	48%	47%	19%

<sup>1</sup> In Dukku two women tested HIV ve+ nine received ARV prophylaxis seven women were to referred to facility reason for 450% is score

<sup>2,3,4</sup> Not colour coded as there are no internationally recognised standards for these indicators.

Data sources: DHIS2

