

What is preterm birth?

A preterm birth is when a baby is born alive but before 37 weeks of pregnancy are completed¹. In Tanzania, one in four newborn deaths are due to preterm birth complications^{4,a}. Babies born too soon are between 6 and 26 times more likely to die during the first four weeks of their lives than babies born at term³.

In addition, preterm survivors often suffer from lifelong disabilities, such as visual and hearing impairments, chronic lung disease, long term cardiovascular ill-health, learning and behavioural impairments¹. Prematurity has a far-reaching impact on their development and on their health as children and adults.

Preterm birth is a public health problem in Tanzania

Tanzania ranks twelfth in the world for the number of preterm births^{2,5,b} - every year 213,500 babies are born before 37 weeks of pregnancy are completed^{2,5,b} - this is one in every 9 babies born alive^{2,5,b}. Prematurity is the second largest cause of death for babies in the first month of life, with 9,400 newborn deaths due to preterm birth every year^{4,c}.

What can be done in Tanzania?

Many of these deaths could be prevented, through both prevention of preterm birth, and care for the premature baby.

- Ⓞ Adolescent pregnancy, older age pregnancy, short time gaps between births, chronic and infectious diseases, and unhealthy pre-pregnancy weight increase the risk of preterm birth¹.
- Ⓞ Family planning strategies, including birth spacing and adolescent-friendly services, can reduce the risk of preterm birth¹.
- Ⓞ Encourage early initiation and exclusive breastfeeding¹.
- Ⓞ Quality antenatal and postnatal care, along with skilled care at delivery, can help to reduce deaths due to preterm birth. Investing in trained and equipped health workers is key, with a focus on the 48 hours surrounding birth¹.
- Ⓞ Newborn resuscitation, antibiotics, hygienic cord care, breastfeeding, and mother to newborn skin-to-skin contact are all simple and effective interventions for saving small and sick babies¹.

Through these largely simple cost-effective solutions and technologies, more than **three quarters of deaths due to prematurity** could be averted with the right care before, during and after delivery.

In 2014 Tanzania committed to the Every Newborn Action Plan⁸ and the Sharpened One Plan⁹. These strategies set out the priority solutions, and call for a united effort to dramatically reduce maternal and newborn deaths, and preventable stillbirths. We must prioritise the evidence-based, cost-effective and feasible solutions in all health facilities that provide pregnancy and delivery services, as well as ensure better data collection on pregnancy outcomes to enable responsive action where it is most needed.

If the Sharpened One Plan is delivered, it could save 9,400 newborn lives and avert 2,500 stillbirths, and prevent 1,400 maternal deaths by December 2015⁵.





Prematurity in Tanzania

Number of newborn deaths in the first month of life (2013)⁶	38,600
Mortality rate in the first month of life (2013) ⁶	21 per 1000 babies born alive
Preterm birth rate (2010) ²	11.4% of babies born alive
Country rank for preterm birth rate (2010)²	70
Number of babies born preterm (2013) ^{2,5,b}	213,500
Country rank for number of babies born preterm (2013)^{2,5,b}	12
Number of deaths due to preterm birth complications (2013) ^{4,c}	9,400
Share of newborn deaths caused by preterm birth complications (2013)^{4,a}	24%
Demand for family planning satisfied (2010) ⁷	61%
Antenatal care of four of more visits (2010)⁷	43%
Skilled attendant at delivery (2010) ⁷	51%
Mothers & babies who received postnatal care within two days of birth (2010)⁷	31%
Early initiation (within one hour) of breastfeeding (2010) ⁷	49%
Exclusive breastfeeding for infants in the first six months (2010)⁷	50%

References:

1. March of Dimes, The Partnership for Maternal Newborn & Child Health, Save the Children, World Health Organization. (2012). *Born Too Soon: The Global Action Report on Preterm Birth*. (Eds) Howson, C.P., Kinney, M.V., & Lawn, J.L. WHO: Geneva.
2. Blencowe, H., Cousens, S., Oestergaard, M. Z., Chou, D., Moller, A.-B., Narwal, R., Adler, A., Garcia, C.V., Rhode, S., Say, L., & Lawn, J. E. (2012). National, regional, and worldwide estimates of preterm birth rates in 2010 with time trends since 1990 for selected countries: a systematic analysis and implications. *The Lancet*, 379(9832), 2162–2172.
3. Katz, J., Lee, A. C. C., Kozuki, N., Lawn, J. E., Cousens, S., Blencowe, H., Ezzati, M., Bhutta, Z., Marchant, T., Willey, B., Adair, L., Barros, F., Baqui, A., Christian, P., Fawzi, W., Gonzalez, R., Humphrey, J., Huybregts, L., Kolsteren, P., Mongkolkeha, A., Mullany, L., Ndyomugenyi, R., Nien, J., Osrin, D., Roberfroid, D., Sania, A., Schmiegelow, C., Silveira, M., Tielsch, J., Vaidya, A., Velaphi, S., Victora, C. G., Watson-Jones, D., & Black, R. E. (2013). Mortality risk in preterm and small-for-gestational-age infants in low-income and middle-income countries: a pooled country analysis. *The Lancet*, 382(9890), 417–25.
4. Liu, L., Oza, S., Hogan, D., Perin, J., Rudan, I., Lawn, J. E., Cousens, S., Mathers, C., & Black, R. E. (2014). Global, regional, and national causes of child mortality in 2000–13, with projections to inform post-2015 priorities: an updated systematic analysis. Supplementary appendix. *The Lancet*, (Early online publication).
5. United Nations Population Division. (forthcoming, 2014). World Population Prospects: The 2013 Revision [estimates for 2013 used].
6. UNICEF. (2014). *Committing to Child Survival: Progress Report 2014 Committing to Child Survival: A Promise Renewed*. New York: United Nations Children's Fund.
7. National Bureau of Statistics [Tanzania], & ICF Macro. (2010). *Tanzania Demographic and Health Survey 2010*. Dar Es Salaam/Calverton: NBS, ICF Macro.
8. World Health Organization, & UNICEF. (2014). *Every Newborn: An Action Plan to End Preventable Deaths*. Geneva: WHO.
9. Ministry of Health and Social Welfare [United Republic of Tanzania]. (2014). *The National Road Map Strategic Plan to Accelerate Reduction of Maternal Newborn and Child Deaths in Tanzania 2008-2015. Sharpened One Plan April 2014*. Dar Es Salaam: MoHSW.

Notes:

- a. In Tanzania, the estimated number of newborn deaths for 2013 is 38,611 and the estimated number of deaths due to preterm birth complications is 9,394.⁴ These estimates were used to calculate the proportion of newborn deaths due to preterm birth complications: $(9,394/38,611) * 100 = 24\%$ (1 in 4).
- b. In Tanzania, the estimated number of live births for 2013 is 1,865,300⁵ and the preterm birth rate for 2010 is 11.44532%². The 2010 preterm birth rate is applied to the 2013 live birth estimate: $(11.44532/100) * 1,865,300 = 213,490$ (rounded to 213,500). The preterm birth rate is 11.4%; this is equivalent to 1 baby out of every 9 being born preterm. Countries are ranked by number of preterm births based on these estimates for 2013.
- c. This is the estimated number of deaths due to preterm birth complications in the first 28 days of life in Tanzania (rounded from 9,394).⁴ If including those who die from preterm birth complications between 0 and 5 years the figure is 10,800 (estimate provided by Save the Children).